

Professor Mark Umstad

MB BS, MD, FRCOG, FRANZCOG
OBSTETRICIAN & GYNAECOLOGIST

MIRENA IUCD : POST-OPERATIVE CARE

General instructions

For the first 24 hours after your procedure you should not work, drive, drink alcohol or operate any heavy machinery. A responsible adult should remain with you overnight.

Eating and drinking

You can eat and drink when you are fully awake after your anaesthetic. Occasionally nausea can be a problem for the first 24 hours but it will settle quickly.

Pain relief

In theatre you will have been given your initial pain relief which is usually a combination of an anti-inflammatory and analgesic. This combination will generally be effective for 6 to 12 hours after your procedure. If you feel any more than mild discomfort following this the best form of treatment is Nurofen tablets. If you have a stomach ulcer or asthma you should not take this medication and try Panadeine as an alternative.

Blood loss

Bleeding for the first 24 to 48 hours after the procedure may be a little heavier than a normal period. The blood loss should then settle quite quickly. Most women have light spotting for up to eight weeks after a Mirena IUCD. This is not unusual. If the bleeding suddenly becomes very heavy or offensive or if you have an associated fever then you should contact me.

Sexual relations

You may resume sexual relations whenever you feel comfortable. This is normally three to seven days after the procedure.

Concerning symptoms

You should notify me if you have **increasing** nausea or vomiting, **increasing** abdominal pain, offensive vaginal bleeding or burning on passing urine.

Review

You should contact the rooms to make an appointment for review approximately eight weeks after your procedure. If you have any concerns or problems before this eight week period do not hesitate to contact me or your general practitioner.

PATIENT COPY